

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.013/ESD.013J Electromagnetics and Applications, Fall 2005

Please use the following citation format:

Markus Zahn, Erich Ippen, and David Staelin, *6.013/ESD.013J Electromagnetics and Applications, Fall 2005*. (Massachusetts Institute of Technology: MIT OpenCourseWare). <http://ocw.mit.edu> (accessed MM DD, YYYY). License: Creative Commons Attribution-Noncommercial-Share Alike.

Note: Please use the actual date you accessed this material in your citation.

For more information about citing these materials or our Terms of Use, visit:  
<http://ocw.mit.edu/terms>

6.013, Electromagnetic Fields, Forces, and Motion  
 Prof. Markus Zahn, Sept. 15, 2005  
**Lecture 3: Differential Form of Maxwell's Equations**

I. Divergence Theorem

1. Divergence Operation




Figure 1-13 The net flux through a closed surface tells us whether there is a source or sink within an enclosed volume.

Courtesy of Krieger Publishing. Used with permission.

$$\oint_S \vec{A} \cdot d\vec{S} = \int_V \text{div}(\vec{A}) dV$$

$$\text{div} \vec{A} = \lim_{\Delta V \rightarrow 0} \frac{\oint_S \vec{A} \cdot d\vec{S}}{\Delta V}$$




Figure 1-14 The flux of a vector  $\mathbf{A}$  through the closed surface  $S$  is given by the surface integral of the component of  $\mathbf{A}$  perpendicular to the surface  $S$ . The differential vector surface area element  $d\mathbf{S}$  is in the direction of the unit normal  $\mathbf{n}$ .

Courtesy of Krieger Publishing. Used with permission.




Figure 1-15 Infinitesimal rectangular volume used to define the divergence of a vector.

Courtesy of Krieger Publishing. Used with permission.

$$\begin{aligned}
\Phi &= \int_1 A_x(x, y, z) dydz - \int_1' A_x(x - \Delta x, y, z) dydz \\
&+ \int_2 A_y(x, y + \Delta y, z) dx dz - \int_2' A_y(x, y, z) dx dz \\
&+ \int_3 A_z(x, y, z + \Delta z) dx dy - \int_3' A_z(x, y, z) dx dy \\
\Phi &\approx \Delta x \Delta y \Delta z \left\{ \frac{[A_x(x, y, z) - A_x(x - \Delta x, y, z)]}{\Delta x} + \frac{[A_y(x, y + \Delta y, z) - A_y(x, y, z)]}{\Delta y} \right. \\
&\quad \left. + \frac{[A_z(x, y, z + \Delta z) - A_z(x, y, z)]}{\Delta z} \right\} \\
&\approx \Delta V \left[ \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right]
\end{aligned}$$

$$\text{div } \bar{A} = \lim_{\Delta V \rightarrow 0} \frac{\oint_S \bar{A} \cdot d\bar{S}}{\Delta V} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}$$

$$\text{Del Operator: } \nabla = \bar{i}_x \frac{\partial}{\partial x} + \bar{i}_y \frac{\partial}{\partial y} + \bar{i}_z \frac{\partial}{\partial z}$$

$$\text{div } \bar{A} = \nabla \cdot \bar{A} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}$$

## 2. Gauss' Integral Theorem





Figure 1-17 Nonzero contributions to the flux of a vector are only obtained across those surfaces that bound the outside of a volume. (a) Within the volume the flux leaving one incremental volume just enters the adjacent volume where (b) the outgoing normals to the common surface separating the volumes are in opposite directions.

Courtesy of Krieger Publishing. Used with permission.

$$\begin{aligned}
\oint_S \bar{A} \cdot d\bar{S} &= \sum_{\substack{i=1 \\ N \rightarrow \infty}}^N \oint_{dS_i} \bar{A} \cdot d\bar{S}_i \\
&= \lim_{\substack{N \rightarrow \infty \\ \Delta V_n \rightarrow 0}} \sum_{i=1}^N (\nabla \cdot \bar{A}) \Delta V_i \\
&= \int_V \nabla \cdot \bar{A} dV \\
\int_V \nabla \cdot \bar{A} dV &= \oint_S \bar{A} \cdot d\bar{a}
\end{aligned}$$



### 3. Gauss' Law in Differential Form

$$\oint_S \epsilon_0 \bar{E} \cdot d\bar{a} = \int_V \nabla \cdot (\epsilon_0 \bar{E}) dV = \int_V \rho dV$$

$$\nabla \cdot (\epsilon_0 \bar{E}) = \rho$$

$$\oint_S \mu_0 \bar{H} \cdot d\bar{a} = \int_V \nabla \cdot (\mu_0 \bar{H}) dV = 0$$


$$\nabla \cdot (\mu_0 \bar{H}) = 0$$

## II. Stokes' Theorem



### 1. Curl Operation

$$\oint_C \bar{A} \cdot d\bar{s} = \int_S \text{Curl}(\bar{A}) \cdot d\bar{a}$$

$$\text{Curl}(\bar{A})_n = \lim_{da_n \rightarrow 0} \frac{\oint_C \bar{A} \cdot d\bar{s}}{da_n}$$



$$\int_S \nabla \times \bar{A} \cdot \bar{da} = \oint_C \bar{A} \cdot \bar{ds}$$



**Figure 1-19** (a) Infinitesimal rectangular contour used to define the circulation. (b) The right-hand rule determines the positive direction perpendicular to a contour.

Courtesy of Krieger Publishing. Used with permission.

$$\begin{aligned}
\oint_C \bar{\mathbf{A}} \cdot \overline{d\mathbf{s}} &= \int_1^{x+\Delta x} A_x(x, y) dx + \int_2^{y+\Delta y} A_y(x + \Delta x, y) dy + \int_3^x A_x(x, y + \Delta y) dx \\
&\quad + \int_4^y A_y(x, y) dy \\
&= \Delta x \Delta y \left[ \frac{[A_x(x, y) - A_x(x, y + \Delta y)]}{\Delta y} + \frac{[A_y(x + \Delta x, y) - A_y(x, y)]}{\Delta x} \right] \\
&= da_z \left[ \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right]
\end{aligned}$$

$$\text{Curl}(\bar{\mathbf{A}})_z = \frac{\oint \bar{\mathbf{A}} \cdot \overline{d\mathbf{s}}}{da_z} = \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y}$$

By symmetry

$$\text{Curl}(\bar{\mathbf{A}})_y = \frac{\oint \bar{\mathbf{A}} \cdot \overline{d\mathbf{s}}}{da_y} = \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x}$$

$$\text{Curl}(\bar{\mathbf{A}})_x = \frac{\oint \bar{\mathbf{A}} \cdot \overline{d\mathbf{s}}}{da_x} = \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z}$$

$$\text{Curl} \bar{\mathbf{A}} = \bar{\mathbf{i}}_x \left[ \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right] + \bar{\mathbf{i}}_y \left[ \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right] + \bar{\mathbf{i}}_z \left[ \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right]$$

$$= \det \begin{bmatrix} \bar{\mathbf{i}}_x & \bar{\mathbf{i}}_y & \bar{\mathbf{i}}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_x & A_y & A_z \end{bmatrix}$$

$$= \nabla \times \bar{\mathbf{A}}$$

## 2. Stokes' Integral Theorem




Figure 1-23 Many incremental line contours distributed over any surface, have nonzero contribution to the circulation only along those parts of the surface on the boundary contour  $L$ .

Courtesy of Krieger Publishing. Used with permission.

$$\begin{aligned} \lim_{N \rightarrow \infty} \sum_{i=1}^N \oint_{dC_i} \bar{A} \cdot d\bar{s}_i &= \oint_C \bar{A} \cdot d\bar{s} \\ &= \sum_{i=1}^{N \rightarrow \infty} (\nabla \times \bar{A}) \cdot d\bar{a}_i \\ &= \int_S (\nabla \times \bar{A}) \cdot d\bar{a} \end{aligned}$$




Figure 1-20 A fluid with a velocity field that has a curl tends to turn the paddle wheel. The curl component found is in the same direction as the thumb when the fingers of the right hand are curled in the direction of rotation.

Courtesy of Krieger Publishing. Used with permission.



### 3. Faraday's Law in Differential Form

$$\oint_c \bar{\mathbf{E}} \cdot d\bar{\mathbf{s}} = \int_s (\nabla \times \bar{\mathbf{E}}) \cdot d\bar{\mathbf{a}} = -\frac{d}{dt} \int_s \mu_0 \bar{\mathbf{H}} \cdot d\bar{\mathbf{a}}$$
$$\nabla \times \bar{\mathbf{E}} = -\mu_0 \frac{\partial \bar{\mathbf{H}}}{\partial t}$$

### 4. Ampère's Law in Differential Form

$$\oint_c \bar{\mathbf{H}} \cdot d\bar{\mathbf{s}} = \int_s \nabla \times \bar{\mathbf{H}} \cdot d\bar{\mathbf{a}} = \int_s \bar{\mathbf{J}} \cdot d\bar{\mathbf{a}} + \frac{d}{dt} \int_s \epsilon_0 \bar{\mathbf{E}} \cdot d\bar{\mathbf{a}}$$
$$\nabla \times \bar{\mathbf{H}} = \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t}$$

## III. Applications to Maxwell's Equations

### 1. Vector Identity

$$\lim_{c \rightarrow 0} \oint_c \bar{\mathbf{A}} \cdot d\bar{\mathbf{s}} = 0 = \oint_s (\nabla \times \bar{\mathbf{A}}) \cdot d\bar{\mathbf{a}} = \int_v \nabla \cdot (\nabla \times \bar{\mathbf{A}}) dV$$

$$\nabla \cdot (\nabla \times \bar{\mathbf{A}}) = 0$$

### 2. Charge Conservation

$$\nabla \cdot \left\{ \nabla \times \bar{\mathbf{H}} = \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t} \right\}$$

$$0 = \nabla \cdot \left[ \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t} \right]$$

$$0 = \nabla \cdot \bar{\mathbf{J}} + \frac{\partial \rho}{\partial t}$$


### 3. Magnetic Field

$$\nabla \cdot \left\{ \nabla \times \bar{\mathbf{E}} = -\mu_0 \frac{\partial \bar{\mathbf{H}}}{\partial t} \right\}$$

$$0 = -\frac{\partial}{\partial t} [\nabla \cdot \mu_0 \bar{\mathbf{H}}] \Rightarrow \nabla \cdot (\mu_0 \bar{\mathbf{H}}) = 0$$

#### 4. Vector Identity

$$\int_a^b \vec{E} \cdot d\vec{l} = \Phi(a) - \Phi(b)$$




if  $a=b$

$$\oint_c \vec{E} \cdot d\vec{l} = \Phi(a) - \Phi(a) = 0$$

$$\vec{E} = -\nabla\Phi$$

$$\oint_c \nabla\Phi \cdot d\vec{l} = 0$$

$$\int_s \nabla \times (\nabla f) \cdot d\vec{a} = \oint_c \nabla f \cdot d\vec{l} = 0 \Rightarrow \nabla \times (\nabla f) = 0$$



IV.

## Summary of Maxwell's Equations in Free Space

Integral Form

Differential Form

Faraday's Law

$$\oint_C \bar{\mathbf{E}} \cdot d\bar{\mathbf{l}} = -\mu_0 \frac{d}{dt} \int_S \bar{\mathbf{H}} \cdot d\bar{\mathbf{a}}$$

$$\nabla \times \bar{\mathbf{E}} = -\mu_0 \frac{\partial \bar{\mathbf{H}}}{\partial t}$$

Ampere's Law

$$\oint_C \bar{\mathbf{H}} \cdot d\bar{\mathbf{l}} = \int_S \bar{\mathbf{J}} \cdot d\bar{\mathbf{a}} + \epsilon_0 \frac{d}{dt} \int_S \bar{\mathbf{E}} \cdot d\bar{\mathbf{a}}$$

$$\nabla \times \bar{\mathbf{H}} = \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t}$$

Gauss' Law

$$\oint_S \epsilon_0 \bar{\mathbf{E}} \cdot d\bar{\mathbf{a}} = \int_V \rho dV$$

$$\nabla \cdot (\epsilon_0 \bar{\mathbf{E}}) = \rho$$

$$\oint_S \mu_0 \bar{\mathbf{H}} \cdot d\bar{\mathbf{a}} = 0$$

$$\nabla \cdot (\mu_0 \bar{\mathbf{H}}) = 0$$

Conservation of charge

$$1. \quad \oint_C \bar{\mathbf{J}} \cdot d\bar{\mathbf{a}} + \frac{d}{dt} \int_V \rho dV = 0$$

$$\nabla \cdot \bar{\mathbf{J}} + \frac{\partial \rho}{\partial t} = 0$$

$$2. \quad \oint_S \left[ \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t} \right] \cdot d\bar{\mathbf{a}} = 0$$

$$\nabla \cdot \left[ \bar{\mathbf{J}} + \epsilon_0 \frac{\partial \bar{\mathbf{E}}}{\partial t} \right] = 0$$

EQS Limit

MQS Limit

$$\nabla \times \bar{\mathbf{E}} \approx 0, \quad \bar{\mathbf{E}} = -\nabla \Phi$$

$$\nabla \times \bar{\mathbf{E}} = -\mu_0 \frac{\partial \bar{\mathbf{H}}}{\partial t}$$

$$\nabla \cdot \bar{\mathbf{E}} = -\nabla \cdot (\nabla \Phi) = -\nabla^2 \Phi = \frac{\rho}{\epsilon_0} \quad (\text{Poisson's Eq.})$$

$$\nabla \times \bar{\mathbf{H}} = \bar{\mathbf{J}}$$

$$\Phi(x, y, z) = \iiint_{x', y', z'} \frac{\rho(x', y', z') dx' dy' dz'}{4\pi\epsilon_0 [(x-x')^2 + (y-y')^2 + (z-z')^2]^{1/2}}$$

$$\nabla \cdot (\mu_0 \bar{\mathbf{H}}) = 0 \Rightarrow \mu_0 \bar{\mathbf{H}} = \nabla \times \bar{\mathbf{A}}$$

$$\nabla^2 \bar{\mathbf{A}} = -\mu_0 \bar{\mathbf{J}}, \quad \nabla \cdot \bar{\mathbf{A}} = 0$$

$$\bar{\mathbf{A}}(x, y, z) = \iiint_{x', y', z'} \frac{\mu_0 \bar{\mathbf{J}}(x', y', z') dx' dy' dz'}{4\pi [(x-x')^2 + (y-y')^2 + (z-z')^2]^{1/2}}$$