

## LECTURE 15

### Poisson process — II

- **Readings:** Finish Section 6.2.
- Review of Poisson process
- Merging and splitting
- Examples
- Random incidence

## Review

- Defining characteristics
  - **Time homogeneity:**  $P(k, \tau)$
  - **Independence**
  - **Small interval probabilities** (small  $\delta$ ):

$$P(k, \delta) \approx \begin{cases} 1 - \lambda\delta, & \text{if } k = 0, \\ \lambda\delta, & \text{if } k = 1, \\ 0, & \text{if } k > 1. \end{cases}$$

- $N_\tau$  is a Poisson r.v., with parameter  $\lambda\tau$ :

$$P(k, \tau) = \frac{(\lambda\tau)^k e^{-\lambda\tau}}{k!}, \quad k = 0, 1, \dots$$

$$\mathbf{E}[N_\tau] = \text{var}(N_\tau) = \lambda\tau$$

- Interarrival times ( $k = 1$ ): exponential:

$$f_{T_1}(t) = \lambda e^{-\lambda t}, \quad t \geq 0, \quad \mathbf{E}[T_1] = 1/\lambda$$

- Time  $Y_k$  to  $k$ th arrival: Erlang( $k$ ):


$$f_{Y_k}(y) = \frac{\lambda^k y^{k-1} e^{-\lambda y}}{(k-1)!}, \quad y \geq 0$$

### Poisson fishing

- Assume: Poisson,  $\lambda = 0.6/\text{hour}$ .
    - Fish for two hours.
    - if no catch, continue until first catch.
- $P(\text{fish for more than two hours}) =$
  - $P(\text{fish for more than two and less than five hours}) =$
  - $P(\text{catch at least two fish}) =$
  - $\mathbf{E}[\text{number of fish}] =$
  - $\mathbf{E}[\text{future fishing time} \mid \text{fished for four hours}] =$
  - $\mathbf{E}[\text{total fishing time}] =$

### Merging Poisson Processes (again)

- Merging of independent Poisson **processes** is Poisson




- What is the probability that the next arrival comes from the first process?

### Light bulb example

- Each light bulb has independent, exponential( $\lambda$ ) lifetime
- Install three light bulbs.  
Find expected time until last light bulb dies out.

### Splitting of Poisson processes


- Assume that email traffic through a server is a Poisson process.  
Destinations of different messages are independent.



- Each output stream is Poisson.

### Random incidence for Poisson

- Poisson process that has been running forever
- Show up at some "random time"  
(really means "arbitrary time")



- What is the distribution of the length of the chosen interarrival interval?

### Random incidence in "renewal processes"

- Series of successive arrivals
  - i.i.d. interarrival times  
(but not necessarily exponential)
- **Example:**  
Bus interarrival times are equally likely to be 5 or 10 minutes
- If you arrive at a "random time":
  - what is the probability that you selected a 5 minute interarrival interval?
  - what is the expected time to next arrival?

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability  
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.