

5A-1 (a) Critical points occur where $x' - y^2 = 0$ and $x - xy = 0$

Now $x' - y^2 = 0 \Rightarrow x = \pm y$

Also $x - xy = 0 \Rightarrow x(1-y) = 0$
 $\Rightarrow x = 0$ or $y = 1$

$\therefore x = 0$ and $y = 0$

OR $y = 1$ and $x = 1$

OR $y = 1$ and $x = -1$

$\therefore (0, 0), (1, 1)$ and $(-1, 1)$ are the critical points

(b) Critical points occur where $1 - x + y = 0$ and $y + 2x^2 = 0$

i.e. $y = x - 1$

Then $0 = x - 1 + 2x^2$

i.e. $x = \frac{1}{2}$ or $x = -1$

But $x = \frac{1}{2} \Rightarrow y = -\frac{1}{2}$

and $x = -1 \Rightarrow y = -2$

$\therefore (\frac{1}{2}, -\frac{1}{2})$ and $(-1, -2)$ are the critical points.

5A-2 (a) Let $y = x'$

Then $y' = x'' = -\mu(x^2 - 1)x' - x$

The autonomous equations are then

$$\begin{cases} x' = y \\ y' = -\mu(x^2 - 1)y - x \end{cases}$$

Critical points occur at

$y = 0$

$-\mu(x^2 - 1)y - x = 0$ i.e. at $(0, 0)$

(b) Let $y = x'$
 Then $y' = x'' = x' - 1 + x^2$

The autonomous equations are then

$$\begin{cases} x' = y \\ y' = y - 1 + x^2 \end{cases}$$

Critical points occur at


$y = 0$

$y - 1 + x^2 = 0 \therefore x^2 = 1 \therefore x = \pm 1$

So the critical points occur


at $(1, 0)$ and $(-1, 0)$

5A-3 (a) For this system the tangent vector $(-f(x, y), -g(x, y))$ to the trajectories is equal in magnitude but opposite in direction to the tangent vector $(f(x, y), g(x, y))$ to the original system. So the trajectories are the same but are traversed in the opposite direction.



The critical points occur at $f(x, y) = 0$ and $y(x, y) = 0$ i.e. the same for both systems.

5A-3 (b) For this system the tangent vector $(g(x, y), -f(x, y))$ to the trajectories is perpendicular to the tangent vector $(f(x, y), g(x, y))$ to the original system. So (b) represents the orthogonal trajectories of the original system.



The critical points of (b) occur at $g(x, y) = 0$ and $-f(x, y) = 0$ i.e. the same as for the original system.

5A-7(a) Let $u = t - t_0$, let $\bar{x}(t) = x_i(t - t_0)$.
Then $x_i(t - t_0) = x_i(u)$ as a function of u
 $= \bar{x}(t)$ as a function of t

[As an example: if $x_i = t^2$, then $x_i(u) = u^2$.
and $\bar{x}(t) = t^2 - 2t_0 t + t_0^2$

By hypothesis: $\frac{dx_i(t)}{dt} = f(x_i(t), y_i(t))$ and $\frac{dx_i(u)}{du} = f(x_i(u), y_i(u))$ (changing letters formally)
 $\frac{dy_i(t)}{dt} = g(x_i(t), y_i(t))$ $\frac{dy_i(u)}{du} = f(x_i(u), y_i(u))$ (*)

But $\frac{d\bar{x}(t)}{dt} = \frac{dx_i(u)}{du} \cdot \frac{du}{dt} = \frac{dx_i(u)}{du}$; similarly $\frac{d\bar{y}(t)}{dt} = \frac{dy_i(u)}{du}$

Therefore, from (*) we get

$\frac{d\bar{x}(t)}{dt} = f(\bar{x}(t), \bar{y}(t))$ which shows that $\bar{x}(t), \bar{y}(t)$ is also a solution.

$\begin{cases} \bar{x}(t) \\ \bar{y}(t) \end{cases} = \begin{cases} x_i(t - t_0) \\ y_i(t - t_0) \end{cases}$ represents the same motion as $\begin{cases} x_i(t) \\ y_i(t) \end{cases}$

but occurring t_0 time-units later.

That is, $\begin{cases} \bar{x}(t, +t_0) \\ \bar{y}(t, +t_0) \end{cases} = \begin{cases} x_i(t) \\ y_i(t) \end{cases}$ so wherever $\begin{cases} x_i \\ y_i \end{cases}$ is at time t , $\begin{cases} \bar{x} \\ \bar{y} \end{cases}$ is there at time $t + t_0$.

[This is the essential property of an autonomous system - the vector field does not change with time, so if we start at a given point t_0 seconds later, we follow the same trajectory path as before, but delayed by t_0 seconds.]

(b) Let $\begin{pmatrix} x_i(t) \\ y_i(t) \end{pmatrix}$ be two trajectories which intersect at (a, b)
i.e. $\begin{pmatrix} x_i(t_0) \\ y_i(t_0) \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} x_i(t_1) \\ y_i(t_1) \end{pmatrix}$ some t_0, t_1 .


By part (a) $\begin{pmatrix} \bar{x}_i(t) \\ \bar{y}_i(t) \end{pmatrix} = \begin{pmatrix} x_i(t - t_0 + t_1) \\ y_i(t - t_0 + t_1) \end{pmatrix}$

is also a solution to the ODE
But $\begin{pmatrix} \bar{x}_i(t_0) \\ \bar{y}_i(t_0) \end{pmatrix} = \begin{pmatrix} x_i(t_1) \\ y_i(t_1) \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$

Thus by the uniqueness theorem $\begin{pmatrix} x_i(t) \\ y_i(t) \end{pmatrix} = \begin{pmatrix} \bar{x}_i(t) \\ \bar{y}_i(t) \end{pmatrix} = \begin{pmatrix} x_i(t - t_0 + t_1) \\ y_i(t - t_0 + t_1) \end{pmatrix}$ for all t

i.e. $\begin{pmatrix} x_i(t) \\ y_i(t) \end{pmatrix}$ and $\begin{pmatrix} x_i(t) \\ y_i(t) \end{pmatrix}$ are the same trajectory and differ at most by a change in parameter.

5B-1 (a) $\frac{y'}{x'} = \frac{dy}{dx} = \frac{-2y}{-x}$



$\frac{dy}{y} = 2 \frac{dx}{x}$
 $\therefore y = Kx^2$

(b) Let $\bar{x}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $M = \begin{pmatrix} -1 & 0 \\ 0 & -2 \end{pmatrix}$

Then $\bar{x}'(t) = M\bar{x}(t)$. This has solution

$\bar{x}(t) = c_1 \bar{v}_1 e^{\lambda_1 t} + c_2 \bar{v}_2 e^{\lambda_2 t}$


where λ_1 and λ_2 are the (distinct) eigenvalues of M with corresponding eigenvectors \bar{v}_1 and \bar{v}_2

Here $\lambda_1 = -1, \lambda_2 = -2$

$\bar{v}_1 = \begin{pmatrix} 1 \\ c \end{pmatrix}, \bar{v}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

Thus $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} c_1 e^{-t} \\ c_2 e^{-2t} \end{pmatrix}$ All trajectories $\rightarrow (0, 0)$ as $t \rightarrow +\infty$

Thus the y phase picture is:



The new trajectories are

$\begin{cases} x = 0 \\ \dot{y} = c_2 e^{-2t} \end{cases}$
($c > 0, < 0, = 0$)


i.e. the positive and negative y -axis, and the trivial trajectory $\bar{x}(t) = 0$ (the origin)

(c) As the picture shows, 3 trajectories are needed to cover a typical solution curve from part (a): λ, λ' , and \bullet (the origin).

(d) This system may be obtained from the original by replacing t by $-t$. Thus we have the same trajectories but with the direction of the arrows reversed.

5B-2

a) $\frac{dy/dt}{dx/dt} = \frac{x}{y} \therefore \frac{dy}{dx} = \frac{x}{y}$ soln: $y^2 - x^2 = c$ hyperbolas shown
 b) $\vec{x} = \begin{bmatrix} x \\ y \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^t + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} e^{-t}$




The only "new" trajectory is the line $\vec{x}(t) = 0$ (the origin)

c) In general, each solution curve (is covered) by one trajectory. However, the two lines $y=x$ and $y=-x$ each require 3 trajectories to cover them.


d) The system $\begin{cases} x' = -y \\ y' = -x \end{cases}$ has the same trajectories as the original system except the arrows are reversed.

5B-3




(a) $\frac{y'}{x'} = \frac{dy}{dx} = \frac{-2x}{y}$
 $y \cdot dy = -2x \cdot dx$
 $\frac{y^2}{2} + x^2 = c$
 These are ellipses for $c > 0$ and the point $(0,0)$ for $c = 0$

(b) For example, along the x-axis ($y=0$), the tangent vectors are at $(x_0, 0)$ is: $\begin{cases} x' = 0 \\ y' = -2x_0 \end{cases}$, i.e., $(0, -2x_0)$



So the direction of motion along the ellipses is clockwise.



5B-4

(a) Let $\vec{x}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $M = \begin{pmatrix} 2 & -3 \\ 1 & -2 \end{pmatrix}$

Then $\vec{x}'(t) = M \vec{x}(t)$

M has eigenvalues $\lambda_1 = 1, \lambda_2 = -1$ with corresponding eigenvectors $\vec{v}_1 = \begin{pmatrix} 3 \\ 1 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$


The system has a critical point at $(0,0)$ which is a saddle point

The general solution is $\vec{x}(t) = c_1 \vec{v}_1 e^{\lambda_1 t} + c_2 \vec{v}_2 e^{\lambda_2 t}$

For $c_1 = 0$ and as $t \rightarrow \infty$
 $\vec{x}(t) = c_2 e^{-t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \rightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

Also for $c_2 = 0$ and $t \rightarrow -\infty$
 $\vec{x}(t) = c_1 e^t \begin{pmatrix} 3 \\ 1 \end{pmatrix} \rightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

Thus the behaviour near the saddle point looks like



(b) Let $\vec{x}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $M = \begin{pmatrix} 2 & 0 \\ 3 & 1 \end{pmatrix}$

Then $\vec{x}'(t) = M \vec{x}(t)$

M has eigenvalues $\lambda_1 = 2, \lambda_2 = 1$

with corresponding eigenvectors $\vec{v}_1 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

The system has an unstable node at $(0,0)$


The general solution is

$\vec{x}(t) = c_1 \begin{pmatrix} 1 \\ 3 \end{pmatrix} e^{2t} + c_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^t$

So as $t \rightarrow \infty$ all trajectories $\rightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

Thus the behaviour near the node looks like:

For $t \approx -\infty, c_1 \begin{pmatrix} 1 \\ 3 \end{pmatrix} e^{2t}$ is dominant term, \therefore solns are near the y-axis
 For $t \approx \infty, c_2 \begin{pmatrix} 0 \\ 1 \end{pmatrix} e^t$ dominates so solns are parallel to $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$



5B-4

(c) Let $\vec{x}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $M = \begin{pmatrix} -2 & -2 \\ -1 & -3 \end{pmatrix}$

Then $\vec{x}'(t) = M\vec{x}(t)$

M has eigenvalues $\lambda_1 = -4, \lambda_2 = -1$
with corresponding eigenvectors $\vec{v}_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \vec{v}_2 = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$
asympt. stable


The system has an node at (0,0)
The general solution is

$$\vec{x}(t) = c_1 \vec{v}_1 e^{-4t} + c_2 \vec{v}_2 e^{-t}$$

As $t \rightarrow \infty$ all trajectories $\rightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

$$\begin{aligned} x(t) &= c_1 e^{-4t} + 2c_2 e^{-t} \\ y(t) &= c_1 e^{-4t} - c_2 e^{-t} \end{aligned}$$

The behavior near the node looks like:
For $t \rightarrow -\infty, (1) e^{-4t}$ dominates so solns are parallel to (1)
For $t \rightarrow \infty, (2) e^{-t}$ dominates, $y=x$ so solns are close to $(\frac{2}{-1})$ "like"




(d) Let $\vec{x}(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $M = \begin{pmatrix} 1 & -2 \\ 1 & 1 \end{pmatrix}$

Then $\vec{x}'(t) = M\vec{x}(t)$

M has eigenvalues $\lambda_1 = 1+i\sqrt{2}, \lambda_2 = 1-i\sqrt{2}$

The system then has an unstable spiral around (0,0).

Then $y=0$
 $x' = x$
 $\therefore x$ is increasing when the spiral cuts the x -axis
As we see e^t behavior the spiral is outward from the origin




$$e) \vec{x}' = \begin{bmatrix} 1 & 1 \\ -2 & -1 \end{bmatrix} \vec{x}$$

Eigenvalues are $\pm i$ (pure imaginary), so the system is a stable center.

(The curves are ellipses, since $\frac{dy}{dx} = \frac{-2x-y}{x+y}$ which integrates easily after cross-multiplying to $2x^2 + 2xy + y^2 = c$)

Direction of motion:

For example, at (1,0), the vector field is $x'=1, y'=-2$



so motion is counterclockwise
(a few other vectors are shown, inaccurately drawn...)

5B-5

(a) Let $y = x'$

Then, assuming $m \neq 0,$

$$y' = x'' = -\frac{c}{m}x' - \frac{k}{m}x$$

The system is then $\begin{cases} x' = y \\ y' = -\frac{k}{m}x - \frac{c}{m}y \end{cases}$

(b) The eigenvalues of $M = \begin{pmatrix} 0 & 1 \\ -\frac{k}{m} & -\frac{c}{m} \end{pmatrix}$


$$\text{are } \lambda_{\pm} = \frac{-c \pm \sqrt{c^2 - 4km}}{2m}$$

(i) $c=0 \Rightarrow \lambda_{\pm} = \pm i\sqrt{\frac{k}{m}}$

Thus there is a stable center at (0,0).

Physically, we'd expect this as putting $c=0$ ($m, k > 0$) in the ODE gives the SHM equation. Thus x and x' are periodic with period $2\pi\sqrt{\frac{m}{k}}$

Thus we expect periodic trajectories in phase space



Here $c^2 - 4km < 0$

(ii) $\sqrt{c^2 - 4km} = 2\sqrt{km} \left(1 - \frac{c^2}{4km}\right)^{1/2}$
or, neglecting $\epsilon,$ $\approx 2\sqrt{km}$

Then $\lambda_{\pm} = -\frac{c}{m} \pm i\sqrt{\frac{k}{m}}$ (eigenvalues)

The behaviour near (0,0) is that of an asymptotically stable spiral (since $-\frac{c}{m} < 0$)

The "radius" of the spiral decays as $t \rightarrow \infty$ like $e^{-\frac{c}{m}t}$ i.e. very slowly indeed!

Physically we have lightly damped harmonic motion e.g. a particle at the end of a spring oscillating in air. The motion is almost simple harmonic but the amplitude of oscillation decays slowly with time.



(iii) No! When $c^2 - 4km \geq 0,$ then as $k, m > 0$

we see $\sqrt{c^2 - 4km} \leq |c|$

Thus adding or subtracting $\sqrt{c^2 - 4km}$ to $-c$ cannot change its sign.

i.e. when the λ 's are real, either they're both positive or both negative. (since $c \geq 0$ always).

5C-5

This one's work, but instructive: think $x' = x - x^2 - xy$ of x, y as 2 population which mutually ~~eat~~ destroy each other: $x - x^2, 3y - 2y^2$ represent their "natural" growth laws, the $-xy$ terms their mutual destruction. [Like two hostile tribes, non-cannibalistic].

5C-1

$x' = x - y + xy$
 $y' = 3x - 2y - xy$

linearization: $x' = x - y$
 $y' = 3x - 2y$
 (at $(0,0)$)

char eqn: $m^2 + m + 1 = 0$
 $m = \frac{-1 \pm \sqrt{-3}}{2}$

\therefore asymp. stable spiral

5C-2

$x' = x + 2x^2 - y^2$
 $y' = x - 2y + x^3$

linear: $x' = x$
 $y' = x - 2y$

eigenvalues are 1, -2 \therefore unstable saddle (since max. is Δ lar)

$\begin{bmatrix} 1 & 0 \\ 1 & -2 \end{bmatrix}$

5C-3

$x' = 2x + y + xy^3$
 $y' = x - 2y - xy$

linear: $x' = 2x + y$
 $y' = x - 2y$

$m^2 - 5 = 0$
 $m = \pm\sqrt{5}$

unstable saddle

$\begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix}$

5C-4

$x' = 1 - y$
 $y' = x^2 - y^2$


critical pts: $1 - y = 0 \Rightarrow y = 1$
 $x^2 - y^2 = 0 \Rightarrow x = \pm 1$ and $(-1, 1)$

in general At $(1, 1)$: since the Jac. matrix (of partial derivs) is $\begin{bmatrix} 0 & -1 \\ 2x & -2y \end{bmatrix}$, the linear is $\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$

$m^2 + 2m + 2 = 0$
 $m = -1 \pm \sqrt{-4} = -1 \pm i$ \therefore asymp. stable spiral

At $(-1, 1)$: linear is (again using Jacobian): $\begin{bmatrix} 0 & -1 \\ -2 & -2 \end{bmatrix}$ $\therefore m^2 + 2m - 2 = 0$
 $m = -1 \pm \sqrt{3}$

\therefore unstable saddle. Eigenvectors: $-\alpha_1, -\alpha_2 = 0$
 $\therefore \begin{bmatrix} 1 \\ -m \end{bmatrix} \therefore \begin{bmatrix} 1 \\ -\sqrt{3} \end{bmatrix}, \begin{bmatrix} 1 \\ 2.73 \end{bmatrix}$



Using this info: (Along dotted line, $y=1$, a few dir. field vectors are drawn, using the original system: $x' = 0$
 $y' = x^2 - 1$)

A few other vectors are drawn in to help the sketch

Critical points: $x(1-x-y) = 0$
 $y(3-2y-x) = 0$

From equation 1, either $x=0$, or $1-x-y=0$.


If $x=0$, eqn 2 says: $y=0$ or $y=3/2$

If $1-x-y=0$, eqn 2 says: either $y=0$ (in which case $1-x=0, x=1$) or $3-2y-x=0$ (in which case we solve the 2 eqns: $1-x-y=0$ getting $y=2$
 $3-2y-x=0$ $x=-1$)

Summary: critical points are $(0,0), (0, 3/2), (1,0), (-1,2)$.

Now we determine their types: Jacobian matrix: $\begin{bmatrix} 1-2x-y & -x \\ x & -x+3-4y \end{bmatrix}$

$(0,0)$: $\begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$ \leftrightarrow unstable node.



the fat lines are impressive pieces of solution curves. Note there is no mutual coexistence! The tribe of always wins, unless there is none of it to start with, essentially because of its stronger growth rate.

5D-1

a) Putting right-side of equations in (2) = 0 gives (assume $x \neq 0, y \neq 0$)

$$-\frac{x}{y} = 1 - x^2 - y^2 = \frac{y}{x} \quad \therefore -x^2 = y^2$$

so $x^2 + y^2 = 0 \quad \therefore \begin{matrix} x=0 \\ y=0 \end{matrix}$
(contradiction)

b) $(\cos t, \sin t)$ satisfies the system (just substitute); trajectory is the unit circle.

c) Equation (3) shows that if $R > 1$, the direction field points in towards the unit \odot , and (along outside of R) if $R < 1$, it points out towards the unit circle. Thus every solution curve is always getting closer to the unit \odot .

5D-2

a) Bendixson criterion:

$$\operatorname{div}(f, g) = (1 + 3x^2) + (1 + 3y^2) > 0$$

\therefore no limit cycle in xy -plane

b) System has no critical points, since $x^2 + y^2 = 0 \Rightarrow x = 0, y = 0$, and this does not make $1 + x - y = 0$.
 \therefore no limit cycles.

c) System has no critical points if $x < -1$, \therefore no limit cycles in this region.

[To see this: $x^2 - y^2 = 0 \Rightarrow y = \pm x$

$$2x + x^2 + y^2 = 0 \Rightarrow 2x + 2x^2 = 0$$

and $y = \pm x \quad \therefore x = 0, -1$

thus critical pts. are $(0, 0), (-1, 1), (-1, -1)$.]

d) Bendixson's criterion:

$$\begin{aligned} \operatorname{div}(f, g) &= a + 2bx - 2cy \\ &\quad + 2cy - 2bx \\ &= a \end{aligned}$$

\therefore no limit cycles if $a \neq 0$.
in xy -plane

5D-3

The system (7) is

$$\begin{aligned} x' &= y \\ y' &= -v(x) - u(x)y \end{aligned}$$

a) By Bendixson's criterion,
 $\operatorname{div}(f, g) = 0 - u(x) < 0$ for all x, y
if $u(x) > 0$.
 \therefore no periodic solution.

b) $v(x) > 0 \Rightarrow$ system has no critical point [at a critical point, $y = 0, \therefore v(x) = 0$]
 \therefore no periodic solution.

5D-5 (like 5D-1)

5E-1 a) linearization is


$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{bmatrix} 1 & -4 \\ 2 & -1 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \text{ at } (0,0).$$

Char. eqn: $\lambda^2 + 7 = 0$
 $(0,0)$ is a center.

For non-lin. system, $(0,0)$ could be a center; or, unstable or asymptotically stable spiral.

b) linearization is

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{bmatrix} 3 & -1 \\ -6 & 2 \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \text{ at } (0,0)$$

char. eqn: $\lambda^2 - 5\lambda = 0, \lambda = 0, 5$
 $\therefore (0,0)$ is not isolated - it is one of a line of critical points,
 all unstable: 

For non-linear system, picture could stay like this; or turn into an unstable node or saddle.

5E-2 a) $x' = y, y' = x(1-x)$ $J = \begin{bmatrix} 0 & 1 \\ 1-2x & 0 \end{bmatrix}$

Crit. pts: $(0,0), (1,0)$
 At $(0,0), J = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \lambda^2 - 1 = 0$
 $\lambda = 1, \vec{v} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}; \lambda = -1, \vec{v} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

This is an unstable saddle.

At $(1,0), J = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \lambda = \pm i$
 This is a center, clockwise motion.

For non-linear system, three possibilities:



5E-2 b) $x' = x^2 - x + y, y' = -y(x^2 + 1)$


Crit. pts: $\begin{cases} x^2 - x - y = 0 \\ -y(x^2 + 1) = 0 \end{cases} \therefore y = 0, x = 0, 1$

Two crit. pts: $(0,0), (1,0)$.


$J = \begin{bmatrix} 2x-1 & 1 \\ -2xy & -x^2-1 \end{bmatrix}$

At $(0,0): J = \begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}, \lambda = -1, \vec{v} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$



repeated incomplete eigenvalue
 asy. stable node



At $(1,0): J = \begin{bmatrix} 1 & 1 \\ 0 & -2 \end{bmatrix}, \lambda_1 = 1, \vec{v}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \lambda_2 = -2, \vec{v}_2 = \begin{pmatrix} -3 \\ -2 \end{pmatrix}$

Picture:  unstable saddle.

For non-linear system, two possibilities:



5E-3 The new system is $x' = \frac{5a}{4}x - px^2, y' = -by + qxy$

whose critical pt is $(\frac{b}{q}, \frac{5a/4}{p})$.

Crit. pt. for the orig. system is: $(\frac{b}{8}, \frac{a}{p})$.

so the effect is to leave the flower population the same, but to increase the beetle population by 25%.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.03 Differential Equations
Spring 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.