

Percentiles of the Normal and t Distributions

	q-0.5	q-0.75	q-0.9	q-0.95	q-0.99	q-0.999
N(0,1)	0.00	0.67	1.28	1.64	2.33	3.09
t (df=1)	0.00	1.00	3.08	6.31	31.82	318.31
t (df=2)	0.00	0.82	1.89	2.92	6.96	22.33
t (df=3)	0.00	0.76	1.64	2.35	4.54	10.21
t (df=4)	0.00	0.74	1.53	2.13	3.75	7.17
t (df=5)	0.00	0.73	1.48	2.02	3.36	5.89
t (df=6)	0.00	0.72	1.44	1.94	3.14	5.21
t (df=7)	0.00	0.71	1.41	1.89	3.00	4.79
t (df=8)	0.00	0.71	1.40	1.86	2.90	4.50
t (df=9)	0.00	0.70	1.38	1.83	2.82	4.30
t (df=10)	0.00	0.70	1.37	1.81	2.76	4.14
t (df=25)	0.00	0.68	1.32	1.71	2.49	3.45
t (df=50)	0.00	0.68	1.30	1.68	2.40	3.26
t (df=100)	0.00	0.68	1.29	1.66	2.36	3.17
t (df=500)	0.00	0.67	1.28	1.65	2.33	3.11

MIT OpenCourseWare
<http://ocw.mit.edu>

18.443 Statistics for Applications
Spring 2015

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.