

18.650
Statistics for Applications

Chapter 5: Parametric hypothesis testing

Cherry Blossom run (1)

- ▶ The credit union Cherry Blossom Run is a 10 mile race that takes place every year in D.C.
- ▶ In 2009 there were 14974 participants
- ▶ Average running time was 103.5 minutes.

Were runners faster in 2012?

To answer this question, select n runners from the 2012 race at random and denote by X_1, \dots, X_n their running time.

Cherry Blossom run (2)

We can see from past data that the running time has Gaussian distribution.

The variance was 373.

Cherry Blossom run (3)

- ▶ We are given i.i.d r.v X_1, \dots, X_n and we want to know if $X_1 \sim \mathcal{N}(103.5, 373)$
- ▶ This is a **hypothesis testing** problem.
- ▶ There are many ways this could be false:
 1. $\mathbb{E}[X_1] \neq 103.5$
 2. $\text{var}[X_1] \neq 373$
 3. X_1 may not even be Gaussian.
- ▶ We are interested in a very specific question: is $\mathbb{E}[X_1] < 103.5$?

Cherry Blossom run (4)

- ▶ We make the following **assumptions**:
 1. $\text{var}[X_1] = 373$ (variance is the same between 2009 and 2012)
 2. X_1 is Gaussian.
- ▶ The only thing that we did not fix is $\mathbb{E}[X_1] = \mu$.
- ▶ Now we want to test (only): “Is $\mu = 103.5$ or is $\mu < 103.5$ ”?
- ▶ By making **modeling assumptions**, we have reduced the number of ways the hypothesis $X_1 \sim \mathcal{N}(103.5, 373)$ may be rejected.
- ▶ The only way it can be rejected is if $X_1 \sim \mathcal{N}(\mu, 373)$ for some $\mu < 103.5$.
- ▶ We compare an expected value to a fixed reference number (103.5).

Cherry Blossom run (5)

Simple heuristic:

“If $\bar{X}_n < 103.5$, then $\mu < 103.5$ ”

This could go wrong if I randomly pick only fast runners in my sample X_1, \dots, X_n .

Better heuristic:

“If $\bar{X}_n < 103.5$ —(something that $\xrightarrow{n \rightarrow \infty} 0$), then $\mu < 103.5$ ”

To make this intuition more precise, we need to take the size of the random fluctuations of \bar{X}_n into account!

Clinical trials (1)

- ▶ Pharmaceutical companies use hypothesis testing to test if a new drug is efficient.
- ▶ To do so, they administer a drug to a group of patients (test group) and a placebo to another group (control group).
- ▶ Assume that the drug is a cough syrup.
- ▶ Let μ_{control} denote the expected number of expectorations per hour after a patient has used the placebo.
- ▶ Let μ_{drug} denote the expected number of expectorations per hour after a patient has used the syrup.
- ▶ We want to know if $\mu_{\text{drug}} < \mu_{\text{control}}$
- ▶ We compare two expected values. No reference number.

Clinical trials (2)

- ▶ Let $X_1, \dots, X_{n_{\text{drug}}}$ denote n_{drug} i.i.d r.v. with distribution $\text{Poiss}(\mu_{\text{drug}})$
- ▶ Let $Y_1, \dots, Y_{n_{\text{control}}}$ denote n_{control} i.i.d r.v. with distribution $\text{Poiss}(\mu_{\text{control}})$
- ▶ We want to test if $\mu_{\text{drug}} < \mu_{\text{control}}$.

Heuristic:

“If $\bar{X}_{\text{drug}} < \bar{X}_{\text{control}} - (\text{something that } \xrightarrow[n_{\text{drug}} \rightarrow \infty, n_{\text{control}} \rightarrow \infty]{} 0)$, then conclude that $\mu_{\text{drug}} < \mu_{\text{control}}$ ”

Heuristics (1)

Example 1: A coin is tossed 80 times, and Heads are obtained 54 times. Can we conclude that the coin is significantly unfair ?

- ▶ $n = 80, X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p)$;
- ▶ $\bar{X}_n = 54/80 = .68$
- ▶ If it was true that $p = .5$: By CLT+Slutsky's theorem,

$$\sqrt{n} \frac{\bar{X}_n - .5}{\sqrt{.5(1 - .5)}} \approx \mathcal{N}(0, 1).$$

- ▶ $\sqrt{n} \frac{\bar{X}_n - .5}{\sqrt{.5(1 - .5)}} \approx 3.22$
- ▶ Conclusion: It **seems quite** reasonable to reject the hypothesis $p = .5$.

Heuristics (2)

Example 2: A coin is tossed 30 times, and Heads are obtained 13 times. Can we conclude that the coin is significantly unfair ?

- ▶ $n = 30, X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p)$;
- ▶ $\bar{X}_n = 13/30 \approx .43$
- ▶ If it was true that $p = .5$: By CLT+Slutsky's theorem,

$$\sqrt{n} \frac{\bar{X}_n - .5}{\sqrt{.5(1 - .5)}} \approx \mathcal{N}(0, 1).$$

- ▶ Our data gives $\sqrt{n} \frac{\bar{X}_n - .5}{\sqrt{.5(1 - .5)}} \approx -.77$
- ▶ The number .77 is a plausible realization of a random variable $Z \sim \mathcal{N}(0, 1)$.
- ▶ Conclusion: our data does not suggest that the coin is unfair.

Statistical formulation (1)

- ▶ Consider a sample X_1, \dots, X_n of i.i.d. random variables and a statistical model $(E, (\mathbb{P}_\theta)_{\theta \in \Theta})$.
- ▶ Let Θ_0 and Θ_1 be disjoint subsets of Θ .
- ▶ Consider the two hypotheses:
$$\begin{cases} H_0 : & \theta \in \Theta_0 \\ H_1 : & \theta \in \Theta_1 \end{cases}$$
- ▶ H_0 is the *null hypothesis*, H_1 is the *alternative hypothesis*.
- ▶ If we believe that the true θ is either in Θ_0 or in Θ_1 , we may want to *test H_0 against H_1* .
- ▶ We want to decide whether to *reject H_0* (look for evidence against H_0 in the data).

Statistical formulation (2)

- ▶ H_0 and H_1 do not play a symmetric role: the data is only used to try to disprove H_0
- ▶ In particular lack of evidence, does not mean that H_0 is true (“innocent until proven guilty”)
- ▶ A *test* is a statistic $\psi \in \{0, 1\}$ such that:
 - ▶ If $\psi = 0$, H_0 is not rejected;
 - ▶ If $\psi = 1$, H_0 is rejected.
- ▶ Coin example: $H_0: p = 1/2$ vs. $H_1: p \neq 1/2$.
- ▶ $\psi = \mathbb{I}\left\{\left|\sqrt{n}\frac{\bar{X}_n - .5}{\sqrt{.5(1 - .5)}}\right| > C\right\}$, for some $C > 0$.
- ▶ How to choose the *threshold* C ?

Statistical formulation (3)

- ▶ *Rejection region* of a test ψ :

$$R_\psi = \{x \in E^n : \psi(x) = 1\}.$$

- ▶ *Type 1 error* of a test ψ (rejecting H_0 when it is actually true):

$$\begin{aligned} \alpha_\psi &: \Theta_0 \rightarrow \mathbb{R} \\ &\theta \mapsto \mathbb{P}_\theta[\psi = 1]. \end{aligned}$$

- ▶ *Type 2 error* of a test ψ (not rejecting H_0 although H_1 is actually true):

$$\begin{aligned} \beta_\psi &: \Theta_1 \rightarrow \mathbb{R} \\ &\theta \mapsto \mathbb{P}_\theta[\psi = 0]. \end{aligned}$$

- ▶ *Power* of a test ψ :

$$\pi_\psi = \inf_{\theta \in \Theta_1} (1 - \beta_\psi(\theta)).$$

Statistical formulation (4)

- ▶ A test ψ has *level* α if

$$\alpha_\psi(\theta) \leq \alpha, \quad \forall \theta \in \Theta_0.$$

- ▶ A test ψ has *asymptotic level* α if

$$\lim_{n \rightarrow \infty} \alpha_\psi(\theta) \leq \alpha, \quad \forall \theta \in \Theta_0.$$

- ▶ In general, a test has the form

$$\psi = \mathbb{I}\{T_n > c\},$$

for some statistic T_n and threshold $c \in \mathbb{R}$.

- ▶ T_n is called the *test statistic*. The rejection region is $R_\psi = \{T_n > c\}$.

Example (1)

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Ber}(p)$, for some unknown $p \in (0, 1)$.
- ▶ We want to test:

$$H_0: p = 1/2 \text{ vs. } H_1: p = 1/2$$

with asymptotic level $\alpha \in (0, 1)$.

- ▶ Let $T_n = \sqrt{n} \frac{\hat{p}_n - 0.5}{\sqrt{.5(1 - .5)}}$, where \hat{p}_n is the MLE.
- ▶ If H_0 is true, then by CLT and Slutsky's theorem,

$$\mathbb{P}[T_n > q_{\alpha/2}] \xrightarrow[n \rightarrow \infty]{} 0.05$$

- ▶ Let $\psi_\alpha = \mathbb{I}\{T_n > q_{\alpha/2}\}$.

Example (2)

Coming back to the two previous coin examples: For $\alpha = 5\%$, $q_{\alpha/2} = 1.96$, so:

- ▶ In **Example 1**, H_0 is rejected at the asymptotic level 5% by the test $\psi_{5\%}$;
- ▶ In **Example 2**, H_0 is not rejected at the asymptotic level 5% by the test $\psi_{5\%}$.

Question: In **Example 1**, for what level α would ψ_α not reject H_0 ? And in **Example 2**, at which level α would ψ_α reject H_0 ?

p-value

Definition

The (asymptotic) *p-value* of a test ψ_α is the smallest (asymptotic) level α at which ψ_α rejects H_0 . It is random, it depends on the sample.

Golden rule

$\text{p-value} \leq \alpha \Leftrightarrow H_0$ is rejected by ψ_α , at the (asymptotic) level α .

The smaller the p-value, the more confidently one can reject H_0 .

- ▶ Example 1: $\text{p-value} = \mathbb{P}[|Z| > 3.21] \ll .01$.
- ▶ Example 2: $\text{p-value} = \mathbb{P}[|Z| > .77] \approx .44$.

Neyman-Pearson's paradigm

Idea: For given hypotheses, among all tests of level/asymptotic level α , is it possible to find one that has maximal power ?

Example: The trivial test $\psi = 0$ that never rejects H_0 has a perfect level ($\alpha = 0$) but poor power ($\pi_\psi = 0$).

Neyman-Pearson's theory provides (the most) powerful tests with given level. In 18.650, we only study several cases.

The χ^2 distributions

Definition

For a positive integer d , the χ^2 (pronounced “Kai-squared”) distribution with d degrees of freedom is the law of the random variable $Z_1^2 + Z_2^2 + \dots + Z_d^2$, where $Z_1, \dots, Z_d \stackrel{iid}{\sim} \mathcal{N}(0, 1)$.

Examples:

▶ If $Z \sim \mathcal{N}_d(\mathbf{0}, I_d)$, then $\|Z\|_2^2 \sim \chi_d^2$.

▶ Recall that the sample variance is given by

$$S_n = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n} \sum_{i=1}^n X_i^2 - (\bar{X}_n)^2$$

▶ Cochran's theorem implies that for $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, if S_n is the sample variance, then

$$\frac{nS_n}{\sigma^2} \sim \chi_{n-1}^2.$$

▶ $\chi_2^2 = \text{Exp}(1/2)$.

Student's T distributions

Definition

For a positive integer d , the *Student's T distribution with d degrees of freedom* (denoted by t_d) is the law of the random variable $\frac{Z}{\sqrt{V/d}}$, where $Z \sim \mathcal{N}(0, 1)$, $V \sim \chi_d^2$ and $Z \perp\!\!\!\perp V$ (Z is independent of V).

Example:

- ▶ Cochran's theorem implies that for $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, if S_n is the sample variance, then

$$\sqrt{n-1} \frac{\bar{X}_n - \mu}{\sqrt{S_n}} \sim t_{n-1}.$$

Wald's test (1)

- ▶ Consider an i.i.d. sample X_1, \dots, X_n with statistical model $(E, (\mathbb{P}_\theta)_{\theta \in \Theta})$, where $\Theta \subseteq \mathbb{R}^d$ ($d \geq 1$) and let $\theta_0 \in \Theta$ be fixed and given.
- ▶ Consider the following hypotheses:

$$\begin{cases} H_0 : \theta = \theta_0 \\ H_1 : \theta \neq \theta_0. \end{cases}$$

- ▶ Let $\hat{\theta}^{MLE}$ be the MLE. Assume the MLE technical conditions are satisfied.
- ▶ If H_0 is true, then

$$\sqrt{n} I(\hat{\theta}^{MLE})^{1/2} \left(\hat{\theta}_n^{MLE} - \theta_0 \right) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_d(0, I_d) \quad \text{w.r.t. } \mathbb{P}_{\theta_0}.$$

Wald's test (2)

- ▶ Hence,

$$\underbrace{n \left(\hat{\theta}_n^{MLE} - \theta_0 \right)^\top I(\hat{\theta}_n^{MLE}) \left(\hat{\theta}_n^{MLE} - \theta_0 \right)}_{T_n} \xrightarrow[n \rightarrow \infty]{(d)} \chi_d^2 \quad \text{w.r.t. } \mathbb{P}_{\theta_0}.$$

- ▶ Wald's test with asymptotic level $\alpha \in (0, 1)$:

$$\psi = \mathbb{I}\{T_n > q_\alpha\},$$

where q_α is the $(1 - \alpha)$ -quantile of χ_d^2 (see tables).

- ▶ Remark: Wald's test is also valid if H_1 has the form “ $\theta > \theta_0$ ” or “ $\theta < \theta_0$ ” or “ $\theta = \theta_1$ ” ...

Likelihood ratio test (1)

- ▶ Consider an i.i.d. sample X_1, \dots, X_n with statistical model $(E, (\mathbb{P}_\theta)_{\theta \in \Theta})$, where $\Theta \subseteq \mathbb{R}^d$ ($d \geq 1$).
- ▶ Suppose the null hypothesis has the form

$$H_0 : (\theta_{r+1}, \dots, \theta_d) = (\theta_{r+1}^{(0)}, \dots, \theta_d^{(0)}),$$

for some fixed and given numbers $\theta_{r+1}^{(0)}, \dots, \theta_d^{(0)}$.

- ▶ Let

$$\hat{\theta}_n = \operatorname{argmax}_{\theta \in \Theta} \ell_n(\theta) \quad (\text{MLE})$$

and

$$\hat{\theta}_n^c = \operatorname{argmax}_{\theta \in \Theta_0} \ell_n(\theta) \quad (\text{"constrained MLE"})$$

Likelihood ratio test (2)

- ▶ Test statistic:

$$T_n = 2 \ell_n(\hat{\theta}_n) - \ell_n(\hat{\theta}_n^c) \quad .$$

- ▶ **Theorem**

Assume H_0 is true and the MLE technical conditions are satisfied.
Then,

$$T_n \xrightarrow[n \rightarrow \infty]{(d)} \chi_{d-r}^2 \quad \text{w.r.t. } \mathbb{P}_\theta.$$

- ▶ Likelihood ratio test with asymptotic level $\alpha \in (0, 1)$:

$$\psi = \mathbb{I}\{T_n > q_\alpha\},$$

where q_α is the $(1 - \alpha)$ -quantile of χ_{d-r}^2 (see tables).

Testing implicit hypotheses (1)

- ▶ Let X_1, \dots, X_n be i.i.d. random variables and let $\theta \in \mathbb{R}^d$ be a parameter associated with the distribution of X_1 (e.g. a moment, the parameter of a statistical model, etc...)
- ▶ Let $g : \mathbb{R}^d \rightarrow \mathbb{R}^k$ be continuously differentiable (with $k < d$).
- ▶ Consider the following hypotheses:

$$\begin{cases} H_0 : g(\theta) = 0 \\ H_1 : g(\theta) \neq 0. \end{cases}$$

- ▶ E.g. $g(\theta) = (\theta_1, \theta_2)$ ($k = 2$), or $g(\theta) = \theta_1 - \theta_2$ ($k = 1$), or...

Testing implicit hypotheses (2)

- ▶ Suppose an asymptotically normal estimator $\hat{\theta}_n$ is available:

$$\sqrt{n} \hat{\theta}_n - \theta \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_d(0, \Sigma(\theta)).$$

- ▶ Delta method:

$$\sqrt{n} g(\hat{\theta}_n) - g(\theta) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_k(0, \Gamma(\theta)),$$

where $\Gamma(\theta) = \nabla g(\theta)^\top \Sigma(\theta) \nabla g(\theta) \in \mathbb{R}^{k \times k}$.

- ▶ Assume $\Sigma(\theta)$ is invertible and $\nabla g(\theta)$ has rank k . So, $\Gamma(\theta)$ is invertible and

$$\sqrt{n} \Gamma(\theta)^{-1/2} g(\hat{\theta}_n) - g(\theta) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_k(0, I_k).$$

Testing implicit hypotheses (3)

- ▶ Then, by Slutsky's theorem, if $\Gamma(\theta)$ is continuous in θ ,

$$\sqrt{n} \Gamma(\hat{\theta}_n)^{-1/2} g(\hat{\theta}_n) - g(\theta) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}_k(0, I_k).$$

- ▶ Hence, if H_0 is true, i.e., $g(\theta) = 0$,

$$\underbrace{ng(\hat{\theta}_n)^\top \Gamma^{-1}(\hat{\theta}_n)g(\hat{\theta}_n)}_{T_n} \xrightarrow[n \rightarrow \infty]{(d)} \chi_k^2.$$

- ▶ Test with asymptotic level α :

$$\psi = \mathbb{I}\{T_n > q_\alpha\},$$

where q_α is the $(1 - \alpha)$ -quantile of χ_k^2 (see tables).

The multinomial case: χ^2 test (1)

Let $E = \{a_1, \dots, a_K\}$ be a finite space and $(\mathbb{P}_{\mathbf{p}})_{\mathbf{p} \in \Delta_K}$ be the family of all probability distributions on E :

$$\blacktriangleright \Delta_K = \left\{ \mathbf{p} = (p_1, \dots, p_K) \in (0, 1)^K : \sum_{j=1}^K p_j = 1 \right\}.$$

\blacktriangleright For $\mathbf{p} \in \Delta_K$ and $X \sim \mathbb{P}_{\mathbf{p}}$,

$$\mathbb{P}_{\mathbf{p}}[X = a_j] = p_j, \quad j = 1, \dots, K.$$

The multinomial case: χ^2 test (2)

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathbb{P}_{\mathbf{p}}$, for some unknown $\mathbf{p} \in \Delta_K$, and let $\mathbf{p}^0 \in \Delta_K$ be fixed.
- ▶ We want to test:

$$H_0: \mathbf{p} = \mathbf{p}^0 \text{ vs. } H_1: \mathbf{p} \neq \mathbf{p}^0$$

with asymptotic level $\alpha \in (0, 1)$.

- ▶ Example: If $\mathbf{p}^0 = (1/K, 1/K, \dots, 1/K)$, we are testing whether $\mathbb{P}_{\mathbf{p}}$ is the uniform distribution on E .

The multinomial case: χ^2 test (3)


- ▶ Likelihood of the model:

$$L_n(X_1, \dots, X_n, \mathbf{p}) = p_1^{N_1} p_2^{N_2} \dots p_K^{N_K},$$

where $N_j = \#\{i = 1, \dots, n : X_i = a_j\}$.

- ▶ Let $\hat{\mathbf{p}}$ be the MLE:

$$\hat{p}_j = \frac{N_j}{n}, \quad j = 1, \dots, K.$$

 $\hat{\mathbf{p}}$ maximizes $\log L_n(X_1, \dots, X_n, \mathbf{p})$ **under the constraint**

$$\sum_{j=1}^K p_j = 1.$$

The multinomial case: χ^2 test (4)

- ▶ If H_0 is true, then $\sqrt{n}(\hat{\mathbf{p}} - \mathbf{p}^0)$ is asymptotically normal, and the following holds.

Theorem

$$\underbrace{n \sum_{j=1}^K \frac{\hat{\mathbf{p}}_j - \mathbf{p}_j^0}{\mathbf{p}_j^0}}_{T_n} \xrightarrow[n \rightarrow \infty]{(d)} \chi_{K-1}^2.$$

- ▶ χ^2 test with asymptotic level α : $\psi_\alpha = \mathbb{I}\{T_n > q_\alpha\}$, where q_α is the $(1 - \alpha)$ -quantile of χ_{K-1}^2 .
- ▶ Asymptotic p -value of this test: p -value = $\mathbb{P}[Z > T_n | T_n]$, where $Z \sim \chi_{K-1}^2$ and $Z \perp\!\!\!\perp T_n$.

The Gaussian case: Student's test (1)

- ▶ Let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma^2)$, for some unknown $\mu \in \mathbb{R}, \sigma^2 > 0$ and let $\mu_0 \in \mathbb{R}$ be fixed, given.
- ▶ We want to test:

$$H_0: \mu = \mu_0 \text{ vs. } H_1: \mu \neq \mu_0$$

with asymptotic level $\alpha \in (0, 1)$.

- ▶ **If σ^2 is known:** Let $T_n = \sqrt{n} \frac{\bar{X}_n - \mu_0}{\sigma}$. Then, $T_n \sim \mathcal{N}(0, 1)$ and

$$\psi_\alpha = \mathbb{I}\{|T_n| > q_{\alpha/2}\}$$

is a test with (non asymptotic) level α .

The Gaussian case: Student's test (2)

If σ^2 is unknown:

- ▶ Let $\widetilde{T}_n = \sqrt{n-1} \frac{\bar{X}_n - \mu_0}{\sqrt{S_n}}$, where S_n is the sample variance.
- ▶ Cochran's theorem:
 - ▶ $\bar{X}_n \perp\!\!\!\perp S_n$;
 - ▶ $\frac{nS_n}{\sigma^2} \sim \chi_{n-1}^2$.
- ▶ Hence, $\widetilde{T}_n \sim t_{n-1}$: Student's distribution with $n - 1$ degrees of freedom.

The Gaussian case: Student's test (3)

- ▶ Student's test with (non asymptotic) level $\alpha \in (0, 1)$:

$$\psi_\alpha = \mathbb{I}\{|\widetilde{T}_n| > q_{\alpha/2}\},$$

where $q_{\alpha/2}$ is the $(1 - \alpha/2)$ -quantile of t_{n-1} .

- ▶ If H_1 is $\mu > \mu_0$, Student's test with level $\alpha \in (0, 1)$ is:

$$\psi'_\alpha = \mathbb{I}\{\widetilde{T}_n > q_\alpha\},$$

where q_α is the $(1 - \alpha)$ -quantile of t_{n-1} .

- ▶ Advantage of Student's test:
 - ▶ Non asymptotic
 - ▶ Can be run on small samples
- ▶ Drawback of Student's test: It relies on the assumption that the sample is Gaussian.

Two-sample test: large sample case (1)

- ▶ Consider two samples: X_1, \dots, X_n and Y_1, \dots, Y_m , of independent random variables such that

$$\mathbb{E}[X_1] = \dots = \mathbb{E}[X_n] = \mu_X$$

, and

$$\mathbb{E}[Y_1] = \dots = \mathbb{E}[Y_m] = \mu_Y$$

- ▶ Assume that the variances of are known so assume (without loss of generality) that

$$\text{var}(X_1) = \dots = \text{var}(X_n) = \text{var}(Y_1) = \dots = \text{var}(Y_m) = 1$$

- ▶ We want to test:

$$H_0: \mu_X = \mu_Y \text{ vs. } H_1: \mu_X \neq \mu_Y$$

with asymptotic level $\alpha \in (0, 1)$.

Two-sample test: large sample case (2)

From CLT:

$$\sqrt{n}(\bar{X}_n - \mu_X) \xrightarrow[n \rightarrow \infty]{(d)} \mathcal{N}(0, 1)$$

and

$$\sqrt{m}(\bar{Y}_m - \mu_Y) \xrightarrow[m \rightarrow \infty]{(d)} \mathcal{N}(0, 1) \quad \Rightarrow \quad \sqrt{n}(\bar{Y}_m - \mu_Y) \xrightarrow[\frac{m}{n} \rightarrow \gamma]{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \mathcal{N}(0, \gamma)$$

Moreover, the two samples are independent so

$$\sqrt{n}(\bar{X}_n - \bar{Y}_m) + \sqrt{n}(\mu_X - \mu_Y) \xrightarrow[\frac{m}{n} \rightarrow \gamma]{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \mathcal{N}(0, 1 + \gamma)$$

Under $H_0 : \mu_X = \mu_Y$:

$$\sqrt{n} \frac{\bar{X}_n - \bar{Y}_m}{\sqrt{1 + m/n}} \xrightarrow[\frac{m}{n} \rightarrow \gamma]{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \mathcal{N}(0, 1)$$

Test:
$$\psi_\alpha = \mathbb{I} \left\{ \sqrt{n} \frac{\bar{X}_n - \bar{Y}_m}{\sqrt{1 + m/n}} > q_{\alpha/2} \right\}$$

Two-sample T-test

- ▶ If the variances are unknown but we know that $X_i \sim \mathcal{N}(\mu_X, \sigma_X^2)$, $Y_i \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$.

- ▶ Then

$$\bar{X}_n - \bar{Y}_m \sim \mathcal{N}\left(\mu_X - \mu_Y, \frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}\right)$$

- ▶ Under H_0 :

$$\frac{\bar{X}_n - \bar{Y}_m}{\sqrt{\sigma_X^2/n + \sigma_Y^2/m}} \sim \mathcal{N}(0, 1)$$

- ▶ For unknown variance:

$$\frac{\bar{X}_n - \bar{Y}_m}{\sqrt{S_X^2/n + S_Y^2/m}} \sim t_N$$

where

$$N = \frac{(S_X^2/n + S_Y^2/m)^2}{\frac{S_X^4}{n^2(n-1)} + \frac{S_Y^4}{m^2(m-1)}}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

18.650 / 18.6501 Statistics for Applications
Fall 2016

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.